
Abuse-Deterrent Opioids — Evaluation and Labeling


Guidance for Industry

**U.S. Department of Health and Human Services
Food and Drug Administration
Center for Drug Evaluation and Research (CDER)**

**Clinical Medical
April 2015**

DEF-00064182

DEF-MDL-10441.00001



Abuse-Deterrent Opioids — Evaluation and Labeling Guidance for Industry

Additional copies are available from:

Office of Communications

Division of Drug Information, WO51, Room 2201

10903 New Hampshire Ave.

Silver Spring, MD 20993-0002

Phone: 301-796-3400; Fax: 301-847-8714

druginfo@fda.hhs.gov

<http://www.fda.gov/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/default.htm>

**U.S. Department of Health and Human Services
Food and Drug Administration
Center for Drug Evaluation and Research (CDER)**

**Clinical Medical
April 2015**

TABLE OF CONTENTS

I.	INTRODUCTION.....	1
II.	BACKGROUND	1
III.	ABUSE-DETERRENT PRODUCTS.....	3
IV.	PREMARKET STUDIES	4
A.	Laboratory Manipulation and Extraction Studies (Category 1)	6
B.	Pharmacokinetic Studies (Category 2).....	8
C.	Clinical Abuse Potential Studies (Category 3)	9
1.	<i>Blinding.....</i>	<i>10</i>
2.	<i>Pre-qualification Phase</i>	<i>10</i>
3.	<i>Assessment Phase</i>	<i>11</i>
4.	<i>Subjects.....</i>	<i>11</i>
5.	<i>Route of Administration, Dose Selection, Manipulation Mode, and Sample Preparation</i>	<i>12</i>
6.	<i>Outcome Measures and Data Interpretation</i>	<i>12</i>
7.	<i>Data Interpretation.....</i>	<i>13</i>
8.	<i>Statistical Analysis.....</i>	<i>14</i>
a.	<i>Background</i>	<i>14</i>
b.	<i>Primary analyses.....</i>	<i>15</i>
c.	<i>Secondary analyses</i>	<i>15</i>
d.	<i>Multiplicity.....</i>	<i>17</i>
V.	POSTMARKET STUDIES (CATEGORY 4)	17
A.	Formal Studies	18
1.	<i>General Characteristics.....</i>	<i>18</i>
2.	<i>Study Design Features.....</i>	<i>19</i>
B.	Supportive Information.....	21
VI.	LABELING	22
VII.	ADDITIONAL RESEARCH NEEDS.....	25

Abuse-Deterrent Opioids — Evaluation and Labeling Guidance for Industry¹

This guidance represents the current thinking of the Food and Drug Administration (FDA or Agency) on this topic. It does not create any rights for any person and is not binding on FDA or the public. You can use an alternative approach if it satisfies the requirements of the applicable statutes and regulations. To discuss an alternative approach, contact the FDA staff responsible for this guidance as listed on the title page.

I. INTRODUCTION

This guidance explains FDA's current thinking about the studies that should be conducted to demonstrate that a given formulation has abuse-deterrent properties. The guidance makes recommendations about how those studies should be performed and evaluated and discusses how to describe those studies and their implications in product labeling.

This guidance is intended to assist sponsors who wish to develop opioid drug products with potentially abuse-deterrent properties and is not intended to apply to products that are not opioids or opioid products that do not have the potential for abuse.

This guidance also does not address issues associated with the development or testing of generic formulations of abuse-deterrent opioid products. FDA intends to address that topic in one or more future guidance documents.

In general, FDA's guidance documents do not establish legally enforceable responsibilities. Instead, guidances describe the Agency's current thinking on a topic and should be viewed only as recommendations, unless specific regulatory or statutory requirements are cited. The use of the word *should* in Agency guidances means that something is suggested or recommended, but not required.

II. BACKGROUND

Prescription opioid products are an important component of modern pain management. However, abuse and misuse of these products have created a serious and growing public health problem. One potentially important step towards the goal of creating safer opioid analgesics has

¹ This guidance has been prepared by the Division of Anesthesia, Analgesia, and Addiction Products, the Office of Regulatory Policy, the Office of Surveillance and Epidemiology, the Office of Biostatistics, and the Controlled Substance Staff in the Center for Drug Evaluation and Research (CDER) at the Food and Drug Administration.

Contains Nonbinding Recommendations

been the development of opioids that are formulated to deter abuse. FDA considers the development of these products a high public health priority.

Because opioid products are often manipulated for purposes of abuse by different routes of administration or to defeat extended-release (ER) properties, most abuse-deterrent technologies developed to date are intended to make manipulation more difficult or to make abuse of the manipulated product less attractive or less rewarding. It should be noted that these technologies have not yet proven successful at deterring the most common form of abuse—swallowing a number of intact capsules or tablets to achieve a feeling of euphoria. Moreover, the fact that a product has abuse-deterrent properties does not mean that there is no risk of abuse. It means, rather, that the risk of abuse is lower than it would be without such properties. Because opioid products must in the end be able to deliver the opioid to the patient, there may always be some abuse of these products.

For purposes of this guidance, *abuse-deterrent properties* are defined as those properties shown to meaningfully **deter** abuse, even if they do not fully **prevent** abuse. The term *abuse* is defined as the intentional, non-therapeutic use of a drug product or substance, even once, to achieve a desirable psychological or physiological effect.² Abuse is not the same as *misuse*, which refers to the intentional therapeutic use of a drug product in an inappropriate way and specifically excludes the definition of abuse.³ This guidance uses the term *abuse-deterrent* rather than *tamper-resistant* because the latter term refers to, or is used in connection with, packaging requirements applicable to certain classes of drugs, devices, and cosmetics.⁴

The science of abuse deterrence is relatively new, and both the formulation technologies and the analytical, clinical, and statistical methods for evaluating those technologies are rapidly evolving. Based on the evolving nature of the field, FDA intends to take a flexible, adaptive approach to the evaluation and labeling of potentially abuse-deterrent products. Methods for evaluating the abuse-deterrent properties of new molecular entities may have to be adapted based on the characteristics of those products and the anticipated routes of abuse. The development of an abuse-deterrent opioid product should be guided by the need to reduce the abuse known or expected to occur with similar products.

Because FDA expects that the market will foster iterative improvements in products with abuse-deterrent properties, no absolute magnitude of effect can be set for establishing abuse-deterrent characteristics. As a result, FDA intends to consider the *totality of the evidence* when reviewing the results of studies evaluating the abuse-deterrent properties of a product.

² Smith S M, Dart R C, Katz N P, et al. 2013. Classification and definition of misuse, abuse, and related events in clinical trials: ACTTION systematic review and recommendations. *Pain*, 154:2287-2296.

³ Ibid.

⁴ FDA's current Good Manufacturing Practice regulations include tamper-evident packaging requirements. See 21 CFR 211.132. There are also requirements for child resistant "special packaging" under the Poison Prevention Packaging Act and regulations adopted by the Consumer Product Safety Commissioner (CPSC) in 16 CFR 1700.

Contains Nonbinding Recommendations

As with all NDA products, FDA intends to consider opioids with abuse-deterrent properties within the context of available therapy. The standard against which each product's abuse-deterrent properties are evaluated will depend on the range of abuse-deterrent and non-abuse-deterrent products on the market at the time of that application.⁵

Abuse-deterrent properties can generally be established only through comparison to another product.

FDA encourages additional scientific and clinical research that will advance the development and assessment of abuse-deterrent technologies.

FDA believes it is critical to address the problem of opioid abuse while seeking to ensure that patients in pain have appropriate access to opioid products. Moreover, it is important that opioids without abuse-deterrent properties remain available for use in some clinical settings. For example, patients in hospice care and with difficulty swallowing may need access to opioid products that are in solution or that can be crushed.

The following section describes the categories of abuse-deterrent products. The premarket and postmarket studies that should be performed to assess the impact of a potentially abuse-deterrent product are discussed in subsequent sections. Finally, information is provided about labeling for abuse-deterrent products.

III. ABUSE-DETERRENT PRODUCTS

Opioid products can be abused in a number of ways. For example, they can be swallowed whole, crushed and swallowed, crushed and snorted, crushed and smoked, or crushed, dissolved and injected. Abuse-deterrent technologies should target known or expected routes of abuse relevant to the proposed product. As a general framework, abuse-deterrent formulations can currently be categorized as follows:

1. *Physical/chemical barriers* – Physical barriers can prevent chewing, crushing, cutting, grating, or grinding of the dosage form. Chemical barriers, such as gelling agents, can resist extraction of the opioid using common solvents like water, simulated biological media, alcohol, or other organic solvents. Physical and chemical barriers can limit drug release following mechanical manipulation, or change the physical form of a drug, rendering it less amenable to abuse.
2. *Agonist/antagonist combinations* – An opioid antagonist can be added to interfere with, reduce, or defeat the euphoria associated with abuse. The antagonist can be sequestered and released only upon manipulation of the product. For example, a drug product can be

⁵ For guidance on the evaluation of abuse potential for purposes of the Controlled Substances Act (CSA), we refer sponsors to FDA's draft guidance for industry *Assessment of Abuse Potential of Drugs*. This guidance is available at: <http://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/UCM198650.pdf>. FDA guidances are available at <http://www.fda.gov/RegulatoryInformation/Guidances/default.htm>.

Contains Nonbinding Recommendations

formulated such that the substance that acts as an antagonist is not clinically active when the product is swallowed, but becomes active if the product is crushed and injected or snorted.

3. *Aversion* – Substances can be added to the product to produce an unpleasant effect if the dosage form is manipulated or is used at a higher dosage than directed. For example, the formulation can include a substance irritating to the nasal mucosa if ground and snorted.
4. *Delivery System* (including use of depot injectable formulations and implants) – Certain drug release designs or the method of drug delivery can offer resistance to abuse. For example, sustained-release depot injectable formulation or a subcutaneous implant may be difficult to manipulate.
5. *New molecular entities and prodrugs*– The properties of a new molecular entity (NME) or prodrug could include the need for enzymatic activation, different receptor binding profiles, slower penetration into the central nervous system, or other novel effects. Prodrugs with abuse-deterrent properties could provide a chemical barrier to the in vitro conversion to the parent opioid, which may deter the abuse of the parent opioid. New molecular entities and prodrugs are subject to evaluation of abuse potential for purposes of the Controlled Substances Act (CSA).
6. *Combination* – Two or more of the above methods could be combined to deter abuse.
7. *Novel approaches* – This category encompasses novel approaches or technologies that are not captured in the previous categories.

IV. PREMARKET STUDIES

First and foremost, any studies designed to evaluate the abuse-deterrent characteristics of an opioid formulation should be scientifically rigorous. Important general considerations for the design of these studies include the appropriateness of positive controls⁶ and comparator drugs, outcome measures, data analyses to permit a meaningful statistical analysis, and selection of subjects for the study.

The evaluation of an abuse-deterrent formulation should take into consideration the known routes of abuse for the non-abuse-deterrent predecessor or similar products, as well as anticipate the effect that deterring abuse by one route may have on shifting abuse to other, possibly riskier route. For example, if a product is known to be abused using nasal and intravenous routes, developing deterrent properties for the nasal route in the absence of deterrent properties for the intravenous route risks shifting abusers from the nasal to the intravenous route, which is associated with a greater risk for the spread of infectious diseases.

Another concept that should be considered is whether the deterrent effects can be expected to have a meaningful impact on the overall abuse of the product. For example, immediate-release (IR) opioid and acetaminophen combination products are predominantly abused using the oral

⁶ For purposes of this guidance, a positive control is an opioid drug product or drug substance expected to result in a predictable opioid drug liking effect and has a known potential for, or history of, abuse.

Contains Nonbinding Recommendations

route. Demonstrating a deterrent effect by the nasal route may not meaningfully reduce overall abuse of the product.

FDA is committed to retaining a flexible, adaptive approach to evaluating potentially abuse-deterrent opioid drug products. This flexibility is intended to permit a sponsor to tailor the development program to suit the abuse-deterrent characteristics of their product and the routes of abuse for that product. The adaptive aspect is intended to permit a sponsor to take into consideration the relevant products on the market at the time they are developing their product, so that appropriate non-abuse-deterrent and abuse-deterrent comparators can be used. For example, for some proposed products the appropriate comparator may be a conventional formulation. However, if there are similar approved products with abuse-deterrent properties described in labeling, the appropriate comparator should be one of those abuse-deterrent products.

The following sections describe three categories of premarket studies. Although, in general, any development program for studying abuse-deterrent technologies should include data from all three categories of studies, there may be exceptions. For example, a formulation with a sequestered antagonist may intentionally be formulated not to resist crushing, so testing the syringeability of the product may not be relevant. In most cases, however, to obtain a full and scientifically rigorous understanding of the impact of a technology or technologies on a product's abuse potential, data from each of the following three categories of premarket studies are appropriate:

1. Laboratory-based in vitro manipulation and extraction studies (Category 1)
2. Pharmacokinetic studies (Category 2)
3. Clinical abuse potential studies (Category 3)

The results of Category 1 studies may influence the design of Category 2 pharmacokinetic studies and Category 3 clinical abuse potential studies by suggesting the methods of manipulation that would yield the greatest release of opioid. The results of Category 2 studies may influence the need for Category 3 studies of clinical abuse potential and the designs and goals of these studies. For example, if the extended-release characteristics of an abuse-deterrent opioid formulation cannot be defeated and the pharmacokinetic profile remains unchanged following oral or nasal administration of the manipulated product, oral and nasal studies of abuse potential may not be necessary.

Additional studies (i.e., Category 4 studies) analyze postmarket data to assess the impact of an abuse-deterrent formulation on actual abuse. Nonclinical drug discrimination studies are useful in the evaluation of the abuse potential of a drug, but their utility in predicting the impact of abuse-deterrent properties on human behavior has not been established.⁷

⁷ See FDA draft guidance for industry, *Assessment of Abuse Potential of Drugs* see <http://www.fda.gov/ucm/groups/fdagov-public/@fdagov-drugs-gen/documents/document/ucm198650.pdf>.

A. Laboratory Manipulation and Extraction Studies (Category 1)

The goal of laboratory-based Category 1 studies should be to evaluate the ease with which the potentially abuse-deterrent properties of a formulation can be defeated or compromised. This information should be used when designing Category 2 and Category 3 studies. These studies are critical to the understanding of product characteristics and performance.⁸

Methodologically, these studies should be designed with knowledge of the physicochemical properties of the product and the methods available to abusers to manipulate the product and should be conducted on the to-be-marketed formulation. Sponsors should consider both the mechanisms by which abusers can be expected to attempt to deliberately overcome the abuse-deterrent properties of the product as well as the ways that patients may alter the formulation (unintentionally or intentionally) that change the rate or amount of drug released (e.g., dose dumping may occur when taking the product with alcohol or when the product is cut, chewed, or crushed). Testing should provide information sufficient to fully characterize the product's abuse-deterrent properties, including the degree of effort required to bypass or defeat those properties. In some cases, when designing in vitro studies, it may be useful to obtain information from prescription opioid abusers about how they would manipulate and abuse an abuse-deterrent product.

In vitro studies should assess various simple and sophisticated mechanical and chemical ways a drug could be manipulated, such as by (1) defeating or compromising the controlled release of an opioid from ER formulations for purposes of abuse by different routes of administration; (2) preparing an IR formulation for alternative routes of administration; or (3) separating the opioid antagonist, if present, from the opioid agonist, thus compromising the product's abuse-deterrent properties. The goal of these studies is to manipulate the product to the point of defeating its abuse-deterrent properties. Once this goal is achieved, it is no longer necessary to continue experiments using more sophisticated methods. For example, if 90% of the opioid can be extracted under a set of conditions in 10 minutes, there is no need to test the same condition for 30 minutes.

The test product should be compared to appropriate comparator products for ease of mechanical manipulation. The ability to crush, cut, grate, or grind the product formulation using readily available items such as spoons, cutters, and coffee grinders should be assessed. Particular attention should be given to particle size distribution following each mode of physical manipulation because particle size may influence the rate of opioid extraction from manipulated product. The effect of heat and cold on mechanical manipulation should also be studied.

Extractability and solubility studies should be designed to determine whether any of the formulation components might be differentially solubilized and extracted, allowing an abuser to

⁸ This topic has been discussed at meetings of the Anesthetic & Life Support Drugs Advisory Committee and the Drug Safety & Risk Management Advisory Committee (*NDA 022272, OxyContin*, May 5, 2008, and September 24, 2009). Additional information on these meetings is available on FDA's web site at the following location: <http://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/Drugs/AnestheticAndAnalgesicDrugProductsAdvisoryCommittee/UCM187082.pdf>.

Contains Nonbinding Recommendations

bypass the drug's abuse-deterrent properties. In addition to extraction and solubility studies, an assessment should be made to determine if free-base opioid can be precipitated from solution by pH adjustment. After establishing how a product could be manipulated, chemical extraction of the opioid from the intact and the manipulated product should be assessed and compared to opioid extraction from the selected intact and similarly manipulated comparator products.

The ease of extracting the opioid from the intact and manipulated product should be determined using a variety of solvents that are commonly available (e.g., water, vinegar, ethanol, isopropanol, acetone, mineral spirits) and those that have potentially relevant solvent characteristics (e.g., pH, polarity, protic vs. aprotic). The effects of time, temperature, pH, and agitation on solvent extraction should also be determined. For products containing more than one drug substance, extractability and solubility studies should be designed to determine whether any of the active ingredients might be differentially solubilized and extracted. Sampling times should start early (e.g., 30 seconds) and continue until at least 80% of the opioid has been released, or 12 hours has been reached. The in vitro drug-release characteristics of the intact and manipulated product should also be compared using a discriminatory and robust dissolution method.

In addition to the general evaluation of the effects of physical and chemical manipulation on the product, there are important route-specific data that should be generated, as follows:

- For a product with potential for abuse by the nasal route, the particle size distribution following attempted manipulation by various methods should be established, and the method that provides the smallest particle size should be used in subsequent studies.
- For a product with potential for abuse by smoking, the amount of drug produced by vaporization at temperatures encompassing the range from the melting point of the active ingredient to its degradation point should be determined. Appropriate controls, such as pure active ingredient, both in salt and free-base form should be included in these assessments.
- For a product with potential for abuse by injection, the amount of opioid that can be obtained in a syringe should be based on studies of intact and manipulated test product and comparator(s) using small volumes of water (5-10 mL) at room temperature and at 90° C – 95° C with and without agitation. Extraction times should range from 30 seconds to 30 minutes. The amount of opioid extracted, the volume of solution collected and the viscosity of the samples should be recorded. The ability to get the sample into a syringe and expel the sample using needles of various gauges should also be explored.

The following examples illustrate the kinds of outcomes that in vitro studies should evaluate.

1. Characteristics of the product by crushing, grinding or melting, or by changing the intact formulation using other methods that would limit nasal administration of the manipulated product, and/or that would limit dissolution of the manipulated product and incorporation into a solvent that could then be injected by intravenous or subcutaneous routes.

Contains Nonbinding Recommendations

2. Quantity of the opioid extracted from the product following the various methods attempted that could be used for injection by intravenous or subcutaneous routes and a description of any barriers resulting from attempts at dissolution for drawing the drug into a syringe.
3. Quantity of opioid antagonist released from an agonist/antagonist combination when it is manipulated for administration by ingestion, nasal administration, or injection.
4. Quantity of opioid product following in vitro manipulation of the prodrug.

B. Pharmacokinetic Studies (Category 2)

The goal of the clinical pharmacokinetic studies, Category 2, should be to understand the in vivo properties of the formulation by comparing the pharmacokinetic profiles of the manipulated formulation with the intact formulation and with manipulated and intact formulations of the comparator drugs through one or more routes of administration. Even though the same routes of administration should be studied for the new product and comparators, if specific circumstances prevent this approach, the study design should be discussed with FDA. The method of manipulation used for the pharmacokinetic studies should be based on the methods explored during in vitro testing that can be expected to result in the greatest drug release. The routes of administration chosen should be relevant to the proposed product, and likely will be based on what is known about the abuse of similar products. Note that, for some development programs, it may be preferable to combine measures of pharmacokinetic parameters for Category 3 studies, in which case separate Category 2 studies may not be necessary.

In general, the pharmacokinetic profile for the oral route of administration should be studied. Appropriate study subjects for Category 2 studies include healthy volunteers as long as naltrexone is used to block the pharmacodynamic effects of the opioids.

Depending on the product, it may be important to evaluate the pharmacokinetic profile for the nasal route of administration as well. For nasal pharmacokinetic studies, it is important to weigh the risk to the subject based on the excipients in the formulation. Only subjects with a history of nasal abuse of opioids should be recruited for these studies. As with the oral route of administration, it may be possible to combine the pharmacokinetic assessment and the pharmacodynamic assessment in one clinical abuse potential study with sampling for the pharmacokinetic analysis.

Relevant pharmacokinetic parameters for the opioid drug and any psychoactive metabolites that should be measured in these studies include the following.

- Maximum concentration ($C_{\max}$)
- Time to maximum concentration ($T_{\max}$)
- Area under the curve (AUC_{0-t} and $AUC_{0-\infty}$)
- Relevant partial AUC, including early time points such as AUC_{0-30} minutes or AUC_{0-2} hours, the period of time when $C_{\max}$ is expected
- Terminal elimination half-life ($T_{1/2}$)

Contains Nonbinding Recommendations

Traditional pharmacokinetic study designs should be employed (e.g., crossover designs), and the results should be analyzed using bioequivalence methods. The rate of rise of drug concentration should be assessed when possible because it is thought to contribute to differential abuse potential among drugs, formulations, and routes of administration.⁹ To support these analyses, it is important to have specimen collection and analysis time points sufficient to cover the onset, peak, and offset of the effects of both IR and ER formulations, in both the intact and manipulated conditions. In addition, these data are necessary to calculate the relevant partial area under the curve, which should capture the time to maximum concentration of the opioid.

If food and alcohol alter the pharmacokinetic parameters of the formulation, data should be provided to characterize those effects.¹⁰ If food significantly increases systemic exposure of the intact formulation, the underlying mechanism for the food effect should be established by assessing whether the effect is based on the drug substance or the formulation and whether the effect is present with intact product as well as with manipulated product. When food is expected to increase exposure, subsequent abuse potential studies of the oral route should be conducted in the fed state to maximize the potential systemic exposure.

In addition to the pharmacokinetic profile of the opioid, for agonist/antagonist combinations, the pharmacokinetic characteristics of the antagonist should be defined for the intact product as well as for the manipulated formulation.

As with all clinical studies, adverse events should be collected, and those that can provide additional insight about the abuse-deterrent effects are especially important. For example, if the manipulated formulation is abused by snorting, it would be important to assess adverse events related to intranasal tolerability.

C. Clinical Abuse Potential Studies (Category 3)

In addition to their use by FDA to formulate its scheduling recommendation under the CSA for drug products containing a controlled substance, clinical studies of abuse potential, Category 3, are important for assessing the impact of potentially abuse-deterrent properties. As discussed in

⁹ References suggesting that drugs associated with a rapid onset of action are associated with greater abuse potential include:

Abreu M E, Bigelow G E, Fleisher L, and Walsh S L. 2001. Effect of intravenous injection speed on responses to cocaine and hydromorphone in humans. *Psychopharmacology*, 154:76-84.

de Wit H, Bodker B, and Ambre J. 1992. Rate of increase of plasma drug level influences subjective responses in humans. *Psychopharmacology*, 107:352-358.

de Wit H, Didish S, and Ambre J. 1993. Subjective and behavioral effects of diazepam depend on its rate of onset. *Psychopharmacology*, 112: 324-330.

¹⁰ FDA has issued a draft guidance on this topic (*Assessment of Abuse Potential of Drugs*). Once finalized, it will represent FDA's current thinking on this topic.

Contains Nonbinding Recommendations

FDA's guidance on that topic,¹¹ the preferred design is a randomized, double-blind, placebo-controlled and positive controlled crossover study. These studies generally are conducted in a drug-experienced, recreational user population. The use of a pre-qualification phase (see section 2 below) to identify subjects who can reproducibly distinguish active drug from placebo is a common enrichment strategy used to improve the power of the study to establish a difference between treatments.

Additional considerations applicable to clinical abuse potential studies used to assess potentially abuse-deterrent properties are discussed below. For products that are not susceptible to manipulation based on Category 1 and 2 testing, study designs for Category 3 testing should be discussed with FDA.

1. Blinding

Clinical studies of abuse potential should use a randomized, double-blind, placebo-controlled and positive controlled crossover design. Because study subjects are recreational drug users and familiar with the effects of the drug substances being studied, the double-dummy technique or other techniques should be used to ensure the blinding of all tests when possible. However, alternative designs may be suitable when the blinding of the study drug and the positive control cannot be maintained and treatment by period interactions may lead to sequence effects in a crossover design. For example, a parallel design may be useful when studying the intranasal route of administration, where subjects may be able to see the differences in volume or color between test drug and placebo or positive control, or when it is not possible to create similar results from manipulation, such as particle size from crushing. In these circumstances, early discussion with FDA is recommended.

For clinical abuse potential studies in which the subjects will snort test samples, administration of the samples in a narrow neck, opaque container with a pre-inserted straw may help facilitate blinding. However, even though subjects might not be able to see the sample, un-blinding may still occur due to the physical properties of samples with similar particle size distribution. In some formulations, higher crushed tablet/capsule volume or larger particle size may inhibit complete intranasal administration thereby contributing to the deterrence effects. To be able to evaluate these effects, it may be necessary to maintain differences in tablet/capsule volume between the potentially abuse-deterrent formulation and the comparator. To facilitate blinding and maintain the crossover design, placebos matched to each of the differing weights or particle sizes may be useful. The details of the preparation of the samples should be provided in the study protocol.

2. Pre-qualification Phase

The purpose of the pre-qualification phase is to increase the power of a study to detect differences in the abuse potential of the various formulations of drug and placebo.¹² In general,

¹¹ Ibid.

¹² An additional advantage of a pre-qualification phase is that it helps familiarize subjects with and train them in the use of various scales and questionnaires that measure subjective effects.

Contains Nonbinding Recommendations

the pre-qualification phase should ensure that subjects can distinguish between placebo and a conventional IR formulation of the same opioid being developed in an abuse-deterrent formulation, using the same route of administration as planned for the assessment phase. There is little value in having subjects unable to distinguish placebo from active drug continue in the study. The positive control should include a strength that is at least equal to the lowest strength selected for the assessment during the clinical phase. An important aspect of the pre-qualification phase is assessing the ability of subjects to tolerate the study dose. If the dose used in the pre-qualification phase is lower than the lowest strength planned for the assessment phase, some subjects may not be able to tolerate the higher dose that will be administered in the assessment phase. Thus, when tolerability may be an issue, particularly if more than one dose is planned for the assessment phase, a pre-qualification dose that is no lower than the lowest dose planned may be the most efficient choice to establish that the subject can distinguish active drug from placebo and can tolerate the study drug in the range to be tested. For example, a 30 mg or 45 mg dose of opioid could be used in the pre-qualification phase when a 30 mg and 60 mg doses will be assessed in the clinical phase.

Qualifying criteria that help identify subjects with an acceptable placebo response and an acceptable response for the positive control should be pre-specified in the study protocol. After a range for an acceptable placebo response is set, a minimum value for the maximum effect ($E_{\max}$) for the positive control should be defined. The minimum $E_{\max}$ for the positive control may vary from measure to measure, and from study to study. However, an acceptable response for the positive control should not overlap with the acceptable range for placebo response.

3. Assessment Phase

The potentially abuse-deterrent product should be compared to a positive control, and the positive control should be compared to placebo to validate the study. For an IR product with potentially abuse-deterrent properties, the positive control should be an IR formulation of the same opioid. For an ER formulation with potentially abuse-deterrent properties, the positive control could be an IR formulation of the same opioid or an ER formulation of the same opioid. In general, these studies should include one strength of the positive control which is associated with high levels of drug liking. However, when assessing drug liking through the intranasal route, the use of two strengths of the positive control may be helpful to both identify a strength of the positive control associated with high drug liking scores and to validate the study.

If there are no approved products with the same drug substance, the positive control should be a drug that, based on pharmacological profile or nonclinical data, can be expected to have similar pharmacodynamic effects. Selection of the positive control in this setting should be discussed with FDA.

4. Subjects

Studies should be conducted in opioid-experienced, recreational drug users who have experience with the particular route of abuse being studied. Subjects should generally not be physically dependent and should not be currently seeking or participating in treatment for drug abuse such that participating in the study could make them vulnerable to relapse. Depending on the

Contains Nonbinding Recommendations

formulation being studied, however, clinical abuse potential studies can be conducted in physically dependent subjects. For example, if the deterrent product contains an opioid antagonist, clinical abuse potential studies in a physically dependent population may provide information not only on the drug liking of the product, but on the ability of the antagonist to precipitate withdrawal in this population.

Detailed characteristics of the study population with respect to past and current drug use and abuse should be captured (e.g., drugs abused, drug of choice, duration of abuse or abstinence).

5. Route of Administration, Dose Selection, Manipulation Mode, and Sample Preparation

The selection of the route(s) of administration should be based on epidemiological data showing that a selected route is a relevant route of abuse. For NMEs, the sponsor should review the relevant routes of abuse for products similar to the test product and discuss the selected routes with FDA. For each relevant route of administration, the potentially abuse-deterrent product and comparator should be manipulated based on the results of Category 1 studies to cause the highest release of the opioid and the highest plasma levels. The dose of the opioid selected for the study should be known to produce high levels of liking in non-tolerant opioid-experienced recreational users.

For studies using the intranasal route of administration, the preparation of the samples is extremely important. The potentially abuse-deterrent product and comparator study drug should be produced with similar particle size distribution based on a detailed protocol for the preparation of the samples, even if different methods are necessary to do so.¹³ With some formulations, a high volume of the crushed tablet/capsule or larger particle size may inhibit complete intranasal administration and, thereby, contribute to deterrence effects. To evaluate these effects, it may be necessary to maintain differences in tablet/capsule volume between the potentially abuse-deterrent product and the comparator.

For studies using the intravenous route of administration, the oral formulations may not be safe for intravenous use depending on the excipients used in the formulation. In place of the manipulated oral formulation, a solution for injection should be prepared using approved, commercially available parenteral products when available, or products suitably formulated for the study. The amount of the opioid and that of the antagonist, when relevant, should be based on extrapolation from in vitro extraction studies of manipulated solid formulations.

6. Outcome Measures and Data Interpretation

In abuse potential studies, the primary method for evaluating the subjective effects of drugs should be through the use of standardized instruments.

¹³ Available safety-related information on the use of the various excipients through the intranasal route should be provided. Additionally, some sponsors have conducted intranasal tolerability studies before the abuse potential studies to evaluate irritation of the nasal cavity, nasal congestion, and discharge, among other measures.

Contains Nonbinding Recommendations

In typical abuse potential studies, several instruments have been used to measure subjective responses predictive of the likelihood of abuse. These instruments include:

- Visual Analogue Scales (VAS) – used for drug liking, good effects, bad effects, and other drug abuse-related effects
- Profile of Mood States

The VAS should be the primary measure for drug liking because it appears to correlate most directly with potential for abuse. Other measures of particular interest include assessment of likelihood to take the drug again and assessment of overall drug liking.¹⁴

These measures can be assessed using either a unipolar or bipolar scale, and a rationale should be provided for the choice for a particular scale. In general, FDA recommends using a bipolar scale for the primary measure of drug liking. Unipolar scales have been used to measure other drug effects, such as good and bad effects. Regardless of whether a unipolar or bipolar scale is selected, FDA recommends that for purposes of training subjects, the same scale be used in the pre-qualification and assessment phases.

7. Data Interpretation

For clinical studies of abuse potential conducted on potentially abuse-deterrent opioid drug products, the primary analysis should be the difference in means of the $E_{\max}$ ¹⁵ for the primary measure(s) based on the population of study completers. A statistical analysis plan (SAP) should be included in the study protocol or submitted as a separate document before un-blinding the study. The sponsor should provide data and dropout information for non-completers. To ensure adequate power, the sponsor should take into account that there will be subjects who drop out of the study early and plan the sample size calculation accordingly. Proper planning should avoid any need to replace subjects who discontinue without completing the study.

Additional pharmacodynamic measures, including positive subjective effects other than drug liking (e.g., take drug again, high, overall drug liking) and other subject-rated assessments, are generally considered secondary endpoints. Other subject-rated assessments of interest include: alertness; drowsiness; nausea; and, when the intranasal route is used, intranasal irritation, burning, need to blow nose, runny nose/nasal discharge, facial pain/pressure, and nasal congestion.

Some sponsors provide descriptive statistics including mean, standard error, median, and interquartile range, calculated for all pharmacodynamic endpoints by time and treatment.¹⁶ What

¹⁴ Overall drug liking measures the user's retrospective assessment of a drug, whereas VAS for drug liking measures the user's immediate assessment.

¹⁵ In general, the primary endpoint of interest is drug liking, and the $E_{\max}$ is captured within 8 hours after dosing. However, the timeframe of measuring the maximum response will be determined by the pharmacokinetic and pharmacodynamic parameters of the formulations studied.

¹⁶ See *Statistical Analysis* Section for further guidance.

Contains Nonbinding Recommendations

constitutes a clinically significant difference in drug liking, between the manipulated and intact versions of the potentially abuse-deterrent product and positive control, is an area requiring further research and will be evaluated on a case-by-case basis. Analysis of postmarket data on abuse levels associated with the potentially abuse-deterrent product being studied may help to support the findings from abuse potential studies.

In addition, when interpreting results from clinical abuse potential studies, attention should be given to the profile of subjective effects produced by the manipulated and intact formulation in terms of onset, peak duration of activity, and offset. The rate of rise of drug onset for the intact and manipulated potentially abuse-deterrent product should be given appropriate weight in the overall analysis of the abuse-deterrent properties. A more rapid onset of action or a shorter time-to-reach peak effect is generally associated with greater abuse potential. Regarding the duration of effect, it may be difficult to interpret the abuse potential of a formulation that produces a sustained liking effect when taken intact or after manipulation, though lower than that produced by the positive control formulation.

The overall assessment of abuse potential should be based on the pattern of findings across all of the measures. In addition, qualitative aspects of the findings, such as the steepness of the drug liking response and duration of the liking effects associated with manipulated formulations, should be taken into consideration, along with other positive effects and negative effects.

8. *Statistical Analysis*

a. Background

The overall goal of a clinical study of abuse potential is to assess a number of abuse potential outcome measures (e.g., drug liking VAS) in the potentially abuse-deterrent product (*T*) relative to a formulation of the drug without abuse-deterrent properties (*C*), or a newly formulated opioid product (positive control). Substantial decreases in the responses for the potentially abuse-deterrent formulation compared to the positive control are evidence of deterrence.

A clinical study of abuse potential should be validated by comparing the responses to *C* with those of placebo (*P*). Thereafter, the assessment of the abuse-deterrence properties of *T* is of primary interest. This can be achieved by comparing the difference in means between *C* and *T* with a *margin* for abuse potential measures and comparing the difference between *C* and *T* relative to *C* in drug liking on a bipolar VAS.

The statistical analysis of the data in a clinical study should begin with descriptive statistics making up tabulations and graphs that include tables of the mean, standard error, and other summary statistics: minimum, Q1, median, Q3, and maximum of the responses of interest for each treatment and for each paired difference among treatments.

Contains Nonbinding Recommendations

Useful graphs include mean time course profiles, heat-maps,¹⁷ and continuous responder profiles.

The next subsection describes the statistical test that sponsors should use for the primary analysis of $E_{\max}$ on the VAS for drug liking. An analysis of the percent reduction in drug liking for T relative to C on the individual level in subsection c is recommended as a secondary analysis.

b. Primary analyses

The primary analysis of abuse-deterrent effects should be based on the comparison of means¹⁸ between crushed, chewed, or otherwise modified T and C with an abuse deterrence margin on drug liking VAS. That is, test

$$H_0 : \mu_C - \mu_T \leq \delta_1 \text{ versus } H_a : \mu_C - \mu_T > \delta_1$$

where $\delta_1 = \delta^*(\mu_C - 50)$, and $0 < \delta^* < 1$. Because C is an opioid drug, the validation test also needs a margin, say δ_2 . That is,

$$H_0 : \mu_C - \mu_P \leq \delta_2 \text{ versus } H_a : \mu_C - \mu_P > \delta_2$$

where $\delta_2 \geq 15$.

The significant level for both tests is 2.5%.

The actual value of δ_1 is related to μ_C , hence, it may vary according to abuse potential measures and the route of drug administration. The δ^* should be pre-specified in the protocol. We also suggest the use of 95% confidence intervals to assess both the differences $\mu_C - \mu_T$ and $\mu_C - \mu_P$.

c. Secondary analyses

In addition to the primary analysis, an analysis should be performed of the percent reduction for the potentially abuse-deterrent product T relative to C from each individual study subject for drug liking VAS on a bipolar scale from 0 to 100. One definition for percent reduction for individual subjects is as follows:

$$\%reduction = \frac{c_i - t_i}{c_i - p_i} \times 100\%, i = 1, 2, \dots, n,$$

where c_i , t_i and p_i are the $E_{\max}$ values for C , T , and P from the i th subject, respectively; n is the sample size.

¹⁷ Chen L and Wang Y. 2012. Heat map displays for data from human abuse potential crossover studies. *Drug Information Journal*, 46:701:707.

¹⁸ If a nonparametric method is necessary, analysis of the median difference in $E_{\max}$ may be appropriate.

Contains Nonbinding Recommendations

However, this definition is problematic because for two subjects having the same $E_{\max}$ values for T and C ($t_1 = t_2$ and $c_1 = c_2$), the larger the placebo response, the greater the percent reduction. A more appropriate definition of percent reduction can be derived by replacing p_i by the neutral score 50 on a bipolar scale; that is,

$$\% \text{ reduction} = \frac{c_i - t_i}{c_i - 50} \times 100\%, i = 1, 2, \dots, n$$

where we assume that $c_i > 50$. In case some subjects have $c_i \leq 50$, define % reduction = 0.

Note that even though most abuse potential studies have a pre-qualification phase, approximately 10% of subjects still have placebo responses p_i over 65, with 5% over 75 in the assessment phase. Consequently, it may be necessary to penalize subjects with large values of p_i in computing percent reduction. For example, the percent reduction could be multiplied by an adjustment factor that equals 1 when p_i is around 50 or less and decreases from 1 when p_i is large. Sponsors should discuss with FDA the need for an adjustment factor in computing percent reduction and an appropriate formula for defining the penalty to be applied before finalizing the study protocol.

Two approaches for assessing the deterrent effects using percent reduction for crossover design studies are provided below. Note that when a parallel design is used, the percent reduction for individual subjects is not applicable, and the primary analysis may also serve the purpose for assessing the percent reduction based on $\mu_C - \mu_T$ related to $\mu_C - 50$.

- Responder Analysis

A *responder* is defined as a subject who had at least $\delta^*100\%$ of reduction, in $E_{\max}$ for T relative to C . To ensure that a majority of subjects are responders, a proportion test can be used to test the null hypothesis that 50% or fewer subjects are responders. That is, test

$$H_0 : p^* \leq 50\% \text{ versus } H_a : p^* > 50\%$$

at the 2.5% significance level where p^* denotes the percentage of responders. The 95% confidence interval of p^* can also be calculated.

- Analysis of the Median Percent Reduction

The median of the percent reduction (ptr) is a descriptive measure of central tendency of ptr . At most 50% of subjects have ptr less than the median, and at most 50% of subjects have ptr greater than the median. If the median of ptr is equal to 30%, for example, it means that approximately 50% of subjects have greater than or equal to a 30% reduction.

Contains Nonbinding Recommendations

For assessing deterrent effects, we can test

$$H_0 : \text{median}(\text{ptr}) \leq \text{DR}\% \text{ versus } H_a : \text{median}(\text{ptr}) > \text{DR}\%$$

at the 2.5% significance level, where DR denotes deterrent reduction. To be consistent with the responder analysis, we recommend $\text{DR}\% = \delta^* 100\%$. If the distribution of *ptr* is symmetric, the Wilcoxon-signed rank test can be used to test the null hypothesis that the $\text{median}(\text{ptr}) \leq \text{DR}\%$, and a 95% confidence interval for the median based on this test can be readily calculated using standard methods. Otherwise, the sign test should be used or an alternate method of this test can be pre-specified in the SAP.

Sponsors should pre-specify one of the two analysis methods for the percent reduction in their SAP in addition to the primary analysis in their clinical studies and discuss with FDA the definition of a responder in the responder analysis or the value of DR% used in the analysis of the median percent reduction before finalizing the study protocol.

d. Multiplicity

Whether or not an adjustment for multiplicity is needed for claiming significant results on the primary or key secondary endpoints varies from study to study. Sponsors should refer to the International Conference on Harmonisation of Technical Requirements for Registration of Pharmaceuticals for Human Use (ICH) guidance *E9 Statistical Principles for Clinical Trials*¹⁹ for statistical principles regarding the multiplicity adjustment.

V. POSTMARKET STUDIES (CATEGORY 4)

Premarket studies focus on assessing the potentially abuse-deterrent properties of a product under controlled conditions. The goal of postmarket²⁰ studies, Category 4, is to determine whether the marketing of a product with abuse-deterrent properties results in meaningful reductions in abuse, misuse, and related adverse clinical outcomes, including addiction, overdose, and death in the post-approval setting. As more abuse-deterrent products are approved, it is possible that the amount of reduction observed in an epidemiologic study may also change. Consequently, a reduction that is deemed meaningful at one time may not be meaningful at another. Given the changing landscape, a numerical threshold cannot define what would be considered a meaningful reduction.

Currently, data on the impact of an abuse-deterrent product on drug abuse in the U.S. population are limited, and thus the optimal data sources, study variables, design features, analytical

¹⁹ ICH guidelines are available on FDA's guidance webpage at <http://www.fda.gov/RegulatoryInformation/Guidances/default.htm>.

²⁰ FDA requires postmarket studies for all opioids with abuse-deterrent labeling claims. For more information on postmarket requirements, see <http://www.fda.gov/Drugs/GuidanceComplianceRegulatoryInformation/Post-marketingPhaseIVCommitments/ucm070766.htm>.

techniques, and outcomes of interest of postmarket epidemiologic studies are not fully established.

Postmarket evaluations of abuse deterrence fall into two categories—formal studies and supportive information. Sponsors should submit protocols to FDA for all formal studies of abuse deterrence. Supportive information can also be submitted to FDA, but cannot substitute for formal studies.

A wide range of interrelated behavioral, clinical, and societal factors contribute to drug abuse; therefore, the effects of an abuse-deterrent formulation can manifest in a variety of ways. Understanding the actual impact of a particular abuse-deterrent formulation may require using a variety of study designs to examine different abuse-related outcomes in given populations of interest. Generally, multiple formal studies using a variety of data sources should be conducted to provide insights into product-specific abuse and the effect of an abuse-deterrent product on the outcomes of interest for other opioid drug products. The use of multiple study designs will also generally help with assessment of the impact of abuse-deterrent products on the full spectrum of abuse-related outcomes (i.e., addiction, overdose, and death) and to characterize and quantify the relevant clinical events that are associated with these outcomes.

Recognizing that the current thinking in this area may change, the following subsections provide recommendations for designing postmarket epidemiologic studies that are capable of detecting a change in the occurrence of abuse as a result of a drug product's abuse-deterrent properties.

A. Formal Studies

1. General Characteristics

Formal studies have the following characteristics:

1. They are hypothesis-driven, population-based, observational evaluations that follow good epidemiological practices^{21,22} and use outcomes that provide meaningful measures of abuse deterrence.
2. They capture one or more outcomes that can be used to assess meaningful reductions in misuse, abuse, addiction, overdose, and death.
3. They produce estimates of abuse and related clinical outcomes that are nationally representative, or are based on data from multiple large geographic regions that can reasonably be generalized to the national level. In the absence of nationally generalizable

²¹ See FDA guidance *Best Practices for Conducting and Reporting Pharmacoepidemiologic Safety Studies Using Electronic Healthcare Data*, available at <http://www.fda.gov/downloads/drugs/guidancecomplianceregulatoryinformation/guidances/ucm243537.pdf>.

²² International Society for Pharmacoepidemiology and Risk Management, *Guidelines for Good Practices and Pharmacoepidemiologic Studies*, available at http://www.pharmacoepi.org/resources/guidelines_08027.cfm, accessed January 25, 2015.

Contains Nonbinding Recommendations

data, smaller or regional studies may be informative, but must be accompanied by a clear explanation of their representativeness and generalizability for appropriate interpretation.

4. They assess overall and route-specific (i.e., injected, snorted, smoked) changes in abuse levels that are associated with an abuse-deterrent product.
5. They are sufficiently powered statistically to assess meaningful changes in drug abuse and are of sufficient duration to examine trends in abuse following the marketing of the abuse-deterrent product. The necessary duration of the studies will depend on a variety of factors, including drug utilization and market share, early postmarket abuse deterrence data, and changes in the prescription opioid or illicit drug market.

2. Study Design Features

The epidemiologic methods and data sources that underlie formal postmarket studies to evaluate the effect of abuse-deterrent formulations are evolving, and best practices have not been established. In addition, characterizing the relevant clinical events that are most useful for understanding the actual impact of a product on abuse-related adverse events is also an evolving science. Based on the current state of this field, we provide below some basic guidelines on recommended study design features that will enable FDA to evaluate the results of formal studies.

1. The study hypothesis and its relationship to assessing abuse deterrence should be clearly stated. The study hypothesis should also include the route(s) of abuse that will be studied.
2. An understanding of each data source is important to the design and interpretation of the study. A description of each data source should be provided in the protocol and should include if and how the data source captures drugs, study outcomes, drug formulation, and route of abuse. The sampling methods, study population, or catchment area for the data source should be clearly described.²³
3. The choice of population(s) in each study should be carefully considered. The populations included in the study should be described in the protocol. At least one study should include a high-risk population, such as a population of known drug abusers, but formal studies should not be limited to only high-risk populations.
4. The protocol and study reports should thoroughly define the study outcomes. The choice of the outcome measure(s) should be justified. Formal studies should, as a group, capture all relevant outcomes: misuse, abuse, addiction, overdose, and death, as well as misuse and abuse clinical outcomes. Overall and route-specific misuse and abuse estimates should include prevalence and frequency of abuse. Clinical outcomes should include, when possible, an assessment of severity of abuse outcomes (e.g., addiction or overdose).

²³ See FDA guidance *Best Practices for Conducting and Reporting Pharmacoepidemiologic Safety Studies Using Electronic Healthcare Data*.

Contains Nonbinding Recommendations

5. Both population- and drug utilization-based estimates should be included in the study protocol.²⁴ Drug utilization-based estimates should use multiple denominators. The denominators are generally the number of prescriptions and the number of extended units (e.g., tablets or capsules). The catchment area for drug utilization data should be specified, particularly for sub-national or regional populations.
6. Sponsors should list all proposed opioid comparators and describe the rationale behind their inclusion. When branded and generic versions of a comparator are marketed, all should be included in the study when possible because many data sources used in abuse studies can identify only active ingredients and do not distinguish between branded and generic products or among multiple generic products. Information should be provided on the ability of data sources and study participants to accurately discriminate among different opioid products and formulations. The choice of comparator is critical for determining if a reduction in drug abuse is the result of a product's abuse-deterrent properties or the result of other factors (e.g., educational programs, prescription drug monitoring programs, changes in law enforcement policies, and the availability of other drugs) or secular trends. The choice of comparators will depend on the particular abuse-deterrent product studied and the opioid market environment at the time the study is initiated. Multiple comparators should be used to achieve the most complete picture of the impact of a product's abuse-deterrent properties. For the purposes of hypotheses, some comparators should be selected and justified as primary comparators in the study protocol before data collection, with additional comparators providing context. The following are examples of several potential abuse-deterrent study comparator scenarios.

If an abuse-deterrent formulation of a previously marketed product is introduced to the market, the primary comparators should include historical and currently available non-abuse-deterrent formulations of the products (including branded and generic whenever possible). Additional individual opioid products should be included as well and should be agreed upon with FDA and identified before the start of the study.

If a new abuse-deterrent product does not have an historical or currently available non-abuse-deterrent version of the same opioid, an appropriate group of comparators should be identified before the start of the study through mutual agreement with FDA. Examples of appropriate primary comparators include immediate release non-abuse-deterrent products with the same active moiety and/or a non-abuse-deterrent product with a relatively stable market share and abuse estimates captured at baseline during the postmarket period. Larger groupings of products can also serve as comparators and can help determine secular trends.

When available, a product that has the same active moiety, but has a different abuse-deterrent property, can serve as a comparator.

²⁴ Secora A, Dormitzer C, Staffa J, and Dal Pan G. 2014. Measures to quantify the abuse of prescription opioids: a review of data sources and metrics. *Pharmacoepidemiology and Drug Safety*, 23(12):1227-37.

Contains Nonbinding Recommendations

7. Understanding the background rates of drug abuse is important for protocol design and interpretation of study results. A baseline assessment of the prevalence of drug abuse for formulations of the same opioid that lack abuse-deterrent properties should be conducted and the baseline time period should be justified.
8. Submissions should include the SAP. The plan should include parameter definitions, unit of analysis, model specification, power and sample size calculations, and any additional variables or predictors. Assessment of the abuse outcome measures should consider both average levels of abuse comparing pre- and post-periods to currently available product (means analysis) and trend analysis.
9. Statistical models should include variables that may affect how the product is used and also other related confounders (e.g., geographic variability and demographic characteristics).
10. Exposure and outcome measures that include self-reported assessments should be validated before the start of the study.
11. The precision of outcome measures will also influence the observational period. Outcome measures with large uncertainty (due to bias or variability) in the exposure or study variable measures, for example, may warrant longer observational periods.
12. Interim analyses are encouraged, but results should be considered tentative in light of their preliminary nature.

B. Supportive Information

Information is considered supportive if it can be used to provide additional context on societal, behavioral, and clinical aspects of abuse and abuse-deterrence. Supportive information may be qualitative or descriptive, and it may rely on sources that capture drug utilization or prescribing patterns, diversion events, attitudes and practices (e.g., tampering) of abusers and other information that may not directly be considered abuse (e.g., data concerning the street value of prescription drugs, information about drug use and misuse from social websites). Investigations that provide supportive information may also include investigations that are conducted in smaller populations or subgroups, and that while perhaps not broadly generalizable, may contribute to the totality of the evidence relating to abuse deterrence.

As is the case for formal studies, best practices for collecting and submitting supportive information are still evolving. However, below are some basic recommendations relating to supportive information.

1. Supportive information should be clearly stated, and the rationale for how the supportive information contributes to a sponsor's portfolio of abuse-related studies should be clearly identified.
2. How supportive information is representative of the population from which it is derived or sampled should be clearly described.

Contains Nonbinding Recommendations

3. How the exposure and outcome are measured should be clearly described along with the relationship between the outcomes measured and the primary outcomes of interest: misuse, abuse, addiction, overdose, and death.
4. Collections of supportive information that include populations of particular interest or geographically diverse settings is strongly encouraged. Overlapping geographic areas between formal and supportive information should be considered.

VI. LABELING

Including information about a product's abuse-deterrent properties in labeling is important to inform health care professionals, the patient community, and the public about a product's abuse potential. Accordingly, FDA encourages sponsors to propose labeling that sets forth the results of in vitro, pharmacokinetic, clinical abuse potential and formal postmarket studies and appropriately characterizes the abuse-deterrent properties of a product.

There are several important concepts about the state of the science of pre- and postmarket studies of abuse deterrence that should be considered as these are reflected in labeling. First, as stated earlier in the guidance, abuse-deterrent does not mean abuse-proof. Therefore, labeling should reflect a product's abuse-deterrent properties, as supported by the data, but should include a caveat that abuse is still possible. Next, premarket studies are intended to demonstrate properties that are predictive of a meaningful abuse-deterrent effect for a particular route of administration. FDA has limited data correlating the abuse-deterrent properties of certain opioid drug products, as demonstrated by premarket studies, with the impact of those properties on abuse or adverse events associated with abuse in the post-approval setting. Even though postmarket studies have the potential to demonstrate such effects, the findings of postmarket studies are not available at the time of initial product approval. Labeling should reflect the predictive quality of premarket studies and include results of relevant completed postmarket studies.

When premarket data show that a product's abuse-deterrent properties can be expected to result in a meaningful reduction in that product's abuse, these data, together with an accurate characterization of what the data mean, should be included in product labeling.²⁵ When postmarket data become available that demonstrate a meaningful reduction in abuse by one or more routes of administration, these data should be added to the product labeling. However, if these postmarket data fail to confirm that the abuse-deterrent properties result in a reduction in abuse, or demonstrate a shift in routes of abuse that represent a greater risk (e.g., a shift from oral and nasal abuse to intravenous abuse), FDA may determine that labeling revisions are needed.

Labeling language regarding abuse deterrence should describe the product's specific abuse-deterrent properties as well as the specific routes of abuse that the product has been developed to deter. For example, a formulation that limits an abuser's ability to crush a tablet and to extract the opioid can be described as limiting manipulation for the purpose of snorting or injection if

²⁵ Abuse-deterrence information in labeling should be presented in the DRUG ABUSE AND DEPENDENCE section under 9.2 Abuse.

Contains Nonbinding Recommendations

the data support such a statement. For this characterization to be accurate and not misleading, however, appropriate caveats are likely to be necessary as described above. For example, a product's labeling should explain that the product's abuse-deterrent properties only make abuse more difficult, not impossible, and that these properties provide no deterrence against other potential forms of abuse.

As noted at the outset of this guidance, FDA will take a flexible, adaptive approach to the evaluation and labeling of abuse-deterrent opioid products. FDA expects sponsors to update their formulations to take advantage of technological improvements and further expects to allow labeling statements related to abuse deterrence commensurate with those advances.

Furthermore, FDA expects sponsors to compare their formulations against approved abuse-deterrent versions of the same opioid. The comparisons should be based on the relevant categories of testing. For instance, if a proposed product is less resistant to manipulation than an approved product, the proposed product may not be eligible for labeling regarding abuse-deterrent properties.

FDA is concerned that, with time, abusers may adapt to abuse-deterrent technologies and discover methods to defeat them. If and when abusers can overcome a technology such that it no longer has a meaningful effect in deterring abuse, FDA may require labeling revisions.

As discussed below, the nature of information in labeling on abuse deterrence for a particular product will depend on the types of studies performed and the result of those studies. Because it cannot provide specific guidance on the magnitude of effect that would be sufficient to support each type of claim, FDA will assess the appropriateness of all proposed labeling statements about abuse deterrence based on the data provided.

Information describing the results of the evaluation of abuse-deterrent properties can be used to support labeling statements based on the three premarket categories (i.e., in vitro data, pharmacokinetic data, and clinical abuse potential studies) and the fourth category (postmarket data) once it is available.

The data necessary to support abuse-deterrent labeling will depend on the characteristics of the product that impart the abuse deterrence and the route of abuse. In general, most abuse-deterrent information included in product labeling will be based on data from more than one category.

Key elements of the study design and conduct should be summarized in the product labeling. Category 1 studies can be described in general terms to avoid creating a *road map* for defeating the product's abuse-deterrent properties. However, the design, conduct, and results of Category 2 and 3 studies should be described in sufficient detail, including the primary outcome measure data from Category 3 studies, to support clear labeling regarding a product's abuse-deterrent properties.

The following are examples of information for inclusion in labeling for different types of abuse-deterrent effects based on various types of premarket studies performed.

Contains Nonbinding Recommendations

- Category 1

For this product, in vitro data demonstrated that an abuse-deterrent product cannot be crushed and dissolved or extracted in a small volume of solution suitable for injection. In this case, Category 1 in vitro data may be sufficient to support a statement in labeling about abuse deterrence for the intravenous route of abuse (See Section IV Premarket Studies). Possible labeling text:

In vitro physical and chemical tablet manipulation studies were performed to evaluate the ability of different extraction methods to defeat the formulation. Results support that Tradename resists crushing, breaking, and dissolution using a variety of tools and solvents and retains some extended-release properties despite manipulation.

These in vitro data demonstrate that Tradename has physical and chemical properties that are expected to deter intravenous abuse. However, abuse of this product is still possible by the oral and nasal routes.

- Category 1 and Category 2

For this product, in vitro and pharmacokinetic data from study of the oral and nasal routes of administration demonstrated that no changes occurred in the extended-release properties of the opioid after crushing or dissolution in a variety of solvents. These data may be sufficient to support statements in labeling about abuse deterrence for the nasal and intravenous routes of abuse. Possible labeling text:

In vitro physical and chemical tablet manipulation studies were performed to evaluate the ability of different extraction methods to defeat the formulation, and pharmacokinetic studies of the oral and intranasal routes were performed to determine the effect of manipulation on drug release. Results support that Tradename resists crushing, breaking, and dissolution using a variety of tools and solvents and retains its extended-release properties despite manipulation.

The in vitro data demonstrate that Tradename has physical and chemical properties that are expected to deter oral, nasal and intravenous abuse. However, abuse of intact product is still possible by the oral route.

- Category 2 and Category 3

For this product, pharmacokinetic and clinical abuse potential studies demonstrated the release of an antagonist from an opioid and antagonist combination product following crushing and that the presence of the antagonist resulted in less drug liking compared to a similar amount of opioid alone when administered by the oral and intranasal routes. In

Contains Nonbinding Recommendations

addition, an additional clinical abuse potential study simulating intravenous abuse using the amounts of opioid and antagonist found to be released from the crushed product also demonstrated reduced drug liking.

The pharmacokinetic data demonstrate that crushing Tradename results in the simultaneous release and rapid absorption of opioid and antagonist. These data along with the results from oral and intranasal clinical abuse potential studies and a clinical abuse potential study of intravenous opioid and antagonist to simulate crushed Tradename indicate that Tradename has properties that are expected to deter abuse via the oral, intranasal, and intravenous routes. However, abuse of Tradename by these routes is still possible.

All of these statements based on Categories 1, 2, or 3 testing should be followed by a statement that data from laboratory and clinical studies may not fully predict abuse potential in the post-approval setting.

As discussed in Section V, postmarket data from a variety of sources can demonstrate that a product's abuse-deterrent properties result in persistent and relevant abuse deterrence. These data can result from appropriately designed, conducted, and analyzed formal postmarket studies and from supportive information on the abuse of the product.

FDA is currently considering formal studies plus a variety of supportive information (e.g., data concerning the street value of prescription drugs) as sources that may be acceptable to provide evidence that a product's formulation has had an actual impact on reducing its abuse. FDA anticipates that data from some or all three of the premarket categories along with data from postmarket studies (including both formal studies and supportive information) would be needed to support a statement in labeling that the product has been shown to reduce abuse. The combined results from all of these studies would be described in the product labeling, including specific study designs, conduct, analyses, and study data.

An example of labeling for a product with evidence of a reduction in abuse is:

These data demonstrated a reduction in the abuse of Tradename in the community setting compared to the levels of abuse, overdose, and death that occurred when only formulations of the same opioid without abuse-deterrent properties were available. This reduction in abuse appears to be attributable to the product's formulation, which deters abuse by injection or snorting of the manipulated product. However, such abuse of this product is still possible, and the product's abuse deterrence properties do not deter abuse associated with swallowing the intact formulation.

VII. ADDITIONAL RESEARCH NEEDS

As discussed above, the science of abuse deterrence is relatively new. Both the technologies involved and the analytical, clinical, and statistical methods for evaluating those technologies are

Contains Nonbinding Recommendations

rapidly evolving. For these reasons, FDA will take a flexible, adaptive approach to the evaluation and labeling of potentially abuse-deterrent opioid products. Additionally, there is considerable room for additional scientific work that could advance the development and assessment of abuse-deterrent products. In particular, FDA encourages additional research on the following topics:

- The quantitative link between changes in the pharmacokinetics of opioids in different formulations and results of a clinical abuse potential study with those same formulations.
- The best assessment methods to employ when analyzing a clinical study of abuse potential.
- The quantitative link between the outcomes from a clinical study of abuse potential comparing formulations and the effect on those same formulations on abuse in the community.
- Further understanding of the best study methods to employ to assess the effect of a product with abuse-deterrent properties on the rates of abuse in the community.
- Development of a communication tool (e.g., a simple graph or chart) to inform prescribers of the relative impact the product has on the different routes of abuse.

Progress on these topics could facilitate the ability of sponsors to propose and FDA to approve labeling that would give a more complete picture of the anticipated effect of products with abuse-deterrent properties. Ultimately, progress in these areas could facilitate product development by reducing the amount of information that is needed to accurately assess a product with abuse-deterrent properties and predict its impact on abuse in the community.