| Please select drug & provide strength (tablets/capsules only): Oxycodone | Patient Name: Please select drug & provide strength (tablets/capsules only): Oxycodone Hydromorphone Methadone Other (optional - district specific) | | |
|---|---|---|---|
| Please select drug & provide strength (tablets/capsules only): Oxycodone | Please select drug & provide strength (tablets/capsules only): Oxycodone | | |
| Check boxes that apply to assist the pharmacist in determining if the prescription should be filled. Attach checklist to hard copy of Rx. Yes No Valid government photo ID copied and attached to hard copy of Rx. For eRx, attach copy at pick- up. ID is optional for Hospice, encology, bedside delivery, sickle cell patients, and patients known 1 to the pharmacy staff, unless it is required by state regulations. No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, 2 prescription must not be dispensed. 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (If cash or a cash discount card, use caution). 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state 7 regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances 8 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriptor's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescriptio | Oxycodone Hydromorphone Methadone Other (optional - district specific) Check boxes that apply to assist the pharmacist in determining if the prescription should be filled. Attach checklist to hard copy of Rx. Valid government photo ID copied and attached to hard copy of Rx. For eRx, attach copy at pick- up. ID is optional for Hospice, oncology, bedside delivery, sickle cell patients, and patients known. 1 to the pharmacy staff, unless it is required by state regulations. No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, 2 prescription must not be dispensed. 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (if cash or a cash discount card, use caution). 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validation makes an expensive the same prescription is being filled on time. If your states tregulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances and be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dose If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients onl | 000000000000000000000000000000000000000 | 000000000000000000000000000000000000000 |
| Check boxes that apply to assist the pharmacist in determining if the prescription should be filled. Attach checklist to hard copy of Rx. Yes No Valid government photo ID copied and attached to hard copy of Rx. For eRx, attach copy at pickup. ID is optional for Hospice, encology, bedside delivery, sickle cell patients, and patients known 1 to the pharmacy staff, unless it is required by state regulations. No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, 2 prescription must not be dispensed. 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ | Check boxes that apply to assist the pharmacist in determining if the prescription should be filled. Attach checklist to hard copy of fix. Valid government photo ID copied and attached to hard copy of Rx. For eRx, attach copy at pickup. ID is optional for Hospice, oncology, bedside delivery, sickle cell patients, and patients known. 1 to the pharmacy staff, unless it is required by state regulations. No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, 2 prescription must not be dispensed. 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (If cash or a cash discount card, use caution). 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validanformation above: Technician Initials Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances and be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dose If in your professional judgment a call to the prescription, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of | | |
| Valid government photo ID copied and attached to hard copy of Rx. For eRx, attach copy at pickup. ID is optional for Hospice, oncology, bedside delivery, sickle cell patients, and patients known to the pharmacy staff, unless it is required by state regulations. No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, prescription must not be dispensed. 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (If cash or a cash discount card, use caution). 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the information above: Technician Initials Technician Initials Steps for Pharmacist to Complete Yes No If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances and be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or glidagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elements of Goo | Valid government photo ID copied and attached to hard copy of Rx. For eRx, attach copy at pickup. ID is optional for Hospice, oncology, bedside delivery, sickle cell patients, and patients known. 1 to the pharmacy staff, unless it is required by state regulations. No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, prescription must not be dispensed. 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (If cash or a cash discount card, use caution). 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validantormation above: Technician Initials Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dose If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: Pharmacist that | | |
| up. ID is optional for Hospice, oncology, bedside delivery, sickle cell patients, and patients known to the pharmacy staff, unless it is required by state regulations. No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, prescription must not be dispensed. Patient has received this prescription from Walgreens before. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is being filled for this previous that the formation above: Technician Initials | up. ID is optional for Hospice, oncology, bedside delivery, sickle cell patients, and patients known. 1 to the pharmacy staff, unless it is required by state regulations. No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, 2 prescription must not be dispensed. 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (If cash or a cash discount card, use caution). 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validation above: Technician Initials Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances and be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or gliagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dose If In your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may lith the prescription without verification provided the of Good Faith Dispensing are met.) | · Yes | Yes No |
| to the pharmacy staff, unless it is required by state regulations. No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, prescription must not be dispensed. Patient has received this prescription from Walgreens before. This prescription is from the same prescriber for the same medication as the previous fill. This prescription is from the same prescriber for the same medication as the previous fill. Patient does not appear intoxicated or under the influence of illicit drugs. Technician Initials Steps for Pharmacist to Complete Yes No If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elements of Good Faith Dispensing are met.) | 1 to the pharmacy staff, unless it is required by state regulations. No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, 2 prescription must not be dispensed. 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (If cash or a cash discount card, use caution). 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validant formation above: Technician Initials Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances an be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or glidagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dose If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: Pharmacist signature Pharmacist signature | - 0 | |
| No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, prescription must not be dispensed. 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (If cash or a cash discount card, use caution). 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the information above: Technician Initials Steps for Pharmacist to Complete Yes No If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient of Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elements of Good Faith Dispensing are met.) | No prior GFD refusal for this exact prescription in patient comments in IC+ profile. If so, prescription must not be dispensed. 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (If cash or a cash discount card, use caution). 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validantormation above: Technician Initials Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances acan be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or glagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dose If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: Patient Laber Drug Good Faith Dispensing Checklist validation procedures and my professional judgment to review this prescription and I have: | <u>n</u> | |
| 2 prescription must not be dispensed. 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (If cash or a cash discount card, use caution). 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the information above: Technician Initials | 2 prescription must not be dispensed. 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (If cash or a cash discount card, use caution). 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validant formation above: Technician Initials Technician Initials Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state 7 regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances 8 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dose If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing ore met.) Notes: attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures and my professional judgment to review this prescription and I have: Dispensed: Pharmacist signature | | |
| 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (If cash or a cash discount card, use caution). 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the information above: Technician Initials Steps for Pharmacist to Complete Yes No If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elements of Good Faith Dispensing are met.) | 3 Patient has received this prescription from Walgreens before. 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (If cash or a cash discount card, use caution). 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validation above: Technician Initials Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dose If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: Patient that I have used the Target Drug Good Faith Dispensing Checklist validation procedures and my professional judgment to review this prescription and I have: Pharmacist signature | | |
| 4 This prescription is from the same prescriber for the same medication as the previous fill. 5 3rd Party Insurance is billed (If cash or a cash discount card, use caution). 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the information above: Technician Initials Steps for Pharmacist to Complete Yes No If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elements of Good Faith Dispensing are met.) | This prescription is from the same prescriber for the same medication as the previous fill. 3rd Party Insurance is billed (If cash or a cash discount card, use caution). 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validation procedures and the validation procedures to validation procedures and the validation procedures and validation procedures and procedures and procedures and my professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures and my professional judgment to review this prescription and I have: Pharmacist signature Pharmacist signature | | |
| 3 3rd Party Insurance is billed (If cash or a cash discount card, use caution). 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the Information above: Technician Initials | 3 and Party Insurance is billed (if cash or a cash discount card, use caution). 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validation above: Technician Initials | | |
| 6 Patient does not appear intoxicated or under the influence of illicit drugs. attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the Information above: Technician Initials | attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validation above: Technician Initials Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosa If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: Pharmacist signature Pharmacist signature Pharmacist signature Pharmacist signature | | |
| attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validate the information above: Technician Initials | attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures to validation procedures. Technician Initials | | |
| Steps for Pharmacist to Complete Steps for Pharmacist to Complete Yes No | Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dose If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: Pharmacist signature Pharmacist signature Pharmacist signature | | |
| Steps for Pharmacist to Complete Yes No | Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state 7 regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances an be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dose If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: Pharmacist signature Pharmacist signature Pharmacist signature | idate th | e the |
| Steps for Pharmacist to Complete Yes No | Steps for Pharmacist to Complete If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dose If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: Pharmacist signature Pharmacist signature | | |
| If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state 7 regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances 8 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elements of Good Faith Dispensing are met.) | If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state 7 regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances 2 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dose If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: Pharmacist signature Pharmacist signature | | |
| If available in your state, PDMP has been reviewed. Prescription is being filled on time. If your state 7 regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances 8 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elements of Good Faith Dispensing are met.) | 7 regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances a can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or gliagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dose If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures and my professional judgment to review this prescription and I have: Dispensed: □ Pharmacist signature □ | Yes | Yes No |
| 7 regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances 8 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elements of Good Faith Dispensing are met.) | 7 regulates early refills of controlled substance prescriptions, follow your state's regulations. Patient and/or prescriber address is within geographical proximity to pharmacy; any variances a can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or gliagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dose If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures and my professional judgment to review this prescription and I have: Dispensed: □ Pharmacist signature □ | | |
| Patient and/or prescriber address is within geographical proximity to pharmacy; any variances 8 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elements of Good Faith Dispensing are met.) | Patient and/or prescriber address is within geographical proximity to pharmacy; any variances 8 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dose If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures and my professional judgment to review this prescription and I have: Pharmacist signature Pharmacist signature | e | |
| 8 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elements of Good Faith Dispensing are met.) | 8 can be reasonably explained. Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dose If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: Pharmacist signature Pharmacist signature | + | - - |
| Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elements of Good Faith Dispensing are met.) | Chronic prescription use can be explained and is supported by documentation (ICD 10 code or 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dose If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures and my professional judgment to review this prescription and I have: Pharmacist signature Pharmacist signature | | |
| 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elements of Good Faith Dispensing are met.) | 9 diagnosis consistent with chronic pain condition). Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dose If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures and my professional judgment to review this prescription and I have: Dispensed: □ Pharmacist signature □ | | |
| Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elements of Good Faith Dispensing are met.) | Per CDC recommendation, naloxone was offered to the patient in case of an emergency for prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dose If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures and my professional judgment to review this prescription and I have: Dispensed: Pharmacist signature | | |
| prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient 10 Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elements of Good Faith Dispensing are met.) | prescriptions ≥ 50 Morphine Milligram Equivalents (MME). *Please refer to the Internal Patient Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dose If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures and my professional judgment to review this prescription and I have: Dispensed: □ Pharmacist signature □ | $+_{\sqcap}$ | ${\Box}$ |
| Talking Points #10-16 Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elements of Good Faith Dispensing are met.) | Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosa If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures and my professional judgment to review this prescription and I have: Dispensed: Pharmacist signature Pharmacist signature | - 1 | |
| Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosage" If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elements of Good Faith Dispensing are met.) | Refer to Opioid Dosage Calculator and CDC handout "Calculating Total Daily Dose of Opioids for Safer Dosa If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures and my professional judgment to review this prescription and I have: Dispensed: Pharmacist signature Pharmacist signature | | |
| If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in notes section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elements of Good Faith Dispensing are met.) | If in your professional judgment a call to the prescriber's clinical staff is warranted, document conversation in section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures and my professional judgment to review this prescription and I have: Dispensed: Pharmacist signature | osage" | |
| section. If no call is required, complete this form with your signature. For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elements of Good Faith Dispensing are met.) | section. If no call is required, complete this form with your signature. (For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures and my professional judgment to review this prescription and I have: Dispensed: Pharmacist signature | | |
| For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the elements of Good Faith Dispensing are met.) | For Hospice, oncology, and sickle cell patients only: RPh may fill the prescription without verification provided the of Good Faith Dispensing are met.) Notes: attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures and my professional judgment to review this prescription and I have: Dispensed: Pharmacist signature | | notes |
| of Good Faith Dispensing are met.) | of Good Faith Dispensing are met.) Notes: attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures and my professional judgment to review this prescription and I have: Dispensed: Pharmacist signature | the elen | elements |
| Notes: | attest that I have used the Target Drug Good Faith Dispensing Checklist validation procedures and my professional judgment to review this prescription and I have: Dispensed: Pharmacist signature | | |
| | orofessional judgment to review this prescription and I have: Dispensed: Pharmacist signature | M0000000000000000000000000000000000000 | 000000000000000000000000000000000000000 |
| | orofessional judgment to review this prescription and I have: Dispensed: Pharmacist signature | | |
| | orofessional judgment to review this prescription and I have: Dispensed: Pharmacist signature | | |
| | orofessional judgment to review this prescription and I have: Dispensed: Pharmacist signature | | |
| | orofessional judgment to review this prescription and I have: Dispensed: Pharmacist signature | | |
| | Dispensed: Pharmacist signature | ıy | |
| | | | |
| | | | |
| professional judgment to review this prescription and I have: | | CSEV | Ev Pur |
| orofessional judgment to review this prescription and I have: Dispensed: Pharmacist signature | , | | |
| orofessional judgment to review this prescription and I have: Dispensed: Pharmacist signature (RPh must inform patient of refusal and make a copy of the Rx for the refusal file folder) et al. 3:18.6 | Proprietary & Confidential. This Checklist and the information contained herein is a Trade Secret of Walgreen Co. | | AG-MD |

WAGMDL00005366

Admitted: 6/02/2022